

Updated study of the quintuple system V994 Her.

P. Zasche¹ and R. Uhlař²

¹ Astronomical Institute, Charles University in Prague, Faculty of Mathematics and Physics, CZ-180 00 Praha 8, V Holešovičkách 2, Czech Republic

² Private Observatory, Pohoří 71, CZ-254 01 Jílové u Prahy, Czech Republic

Received April 6, 2016

ABSTRACT

Aims. We present a new updated study of the unique quintuple system of two eclipsing binaries, V994 Her.

Methods. Based on new obtained photometric observations, we analysed the period variations of the two eclipsing pairs of V994 Her.

Results. We found that the orbital period of the two eclipsing binaries around a common barycenter is about 2.9 years instead of the original period, which was reported as 6.3 years. Moreover, the system now seems to be close to coplanar ($i \approx 84^\circ$), and we also revised the individual masses of all four eclipsing components. The two eclipsing pairs show slow apsidal motion with periods of about 116 and 111 years, respectively. Pairs A and B are distant by about 17 mas on the sky, which is encouraging for observers because an interferometric detection may be possible in the next years.

Conclusions. All of these results are based on the photometric observations of V994 Her and its detailed analysis of the period changes of the two eclipsing binaries.

Key words. stars: binaries: eclipsing – binaries: visual – stars: fundamental parameters – stars: individual: V994 Her.

1. Introduction

The eclipsing system V994 Her was classified as an eclipsing binary of Algol type by Kazarovets et al. (1999), who also published a short remark about a possible variability of the secondary component. Even though the system was observed by the Hipparcos satellite (Perryman et al., 1997), its true nature was not discovered until the detailed study of V994 Her by Lee et al. (2008). Its quadruple nature of two eclipsing binaries was recognized, which was the first discovery of such a system in the whole sky. The system still remains interesting today because the two eclipsing pairs are being monitored for more than 20 years now, and the orbital evolution of both binaries can be studied.

Quadruple (or even higher-order multiple) systems like V994 Her can teach us to better understand the formation and evolution of stars and stellar systems. They may answer questions like how such complex structures were formed (Tohline, 2002), how they evolve into their current state (Tokovinin, 2008), what the preferred initial configuration of the system parameters is, and whether we can trace it somehow from the current configuration (Goodwin & Kroupa, 2005), or we may find out the multiplicity fraction of these stars (Duchêne & Kraus, 2013). All of these interesting questions can partly be answered through a detailed study of eclipsing binaries as parts of complex multiple systems. Following the theoretical modelling, for example, indicates that the so-called Heggie-Hills law plays a role (Heggie, 1975) for a tight close binary and a distant companion, but also for a configuration where the mass ratio of the close pair tends to be closer to unity (see e.g. Goodwin et al. 2007). These two aspects can clearly be

seen in the system V994 Her. Even the existence of planets in such a system cannot be easily ruled out. We know of several planetary systems that orbit eclipsing binaries, for instance, Kepler 47 (Orosz et al., 2012).

About three years ago, we published (Zasche & Uhlař, 2013) the first study of the orbital motion of the system V994 Her based on new eclipse-timing observations. We found that the system is the only one known complex system of two eclipsing binaries orbiting a common barycenter on a rather eccentric orbit ($e=0.747$) and with the orbital period of about 6.3 years. Despite its rather poor coverage in the $O-C$ diagrams, we were able to identify the (rather sharp) period changes of the two pairs around a periastron. From this solution it followed that the two pairs orbit each other on a slightly inclined orbit with an inclination of about 37° .

2. Inconsistency of our original solution for V994 Her

The double eclipsing system V994 Her was first correctly separated into two independent periods in 2008 by Lee et al. (2008). The authors found that the orbital periods of the two pairs are of about $P_A = 2.08$ days, and $P_B = 1.42$ days. Using detailed spectroscopic observations, they were also able to derive the individual masses of the eclipsing components and their spectral types: pair A (B8V + A0V) and pair B (A2V + A4V). All four components are apparently clearly detached and are still placed on the main sequence. Another more distant (about $1''$) component was discovered and is listed in the Washington Double Star Catalog (Mason et al., 2001), which means that this is a quintuple stellar system. At the time of Lee et al. publication, it was the only known such system with two eclipsing

Send offprint requests to: Petr Zasche, e-mail: zasche@sirrah.troja.mff.cuni.cz

binaries located on the sky. The number of such systems has increased today, mainly as a result of the OGLE detections of the Magellanic Cloud eclipsing binaries. For example, Pawlak et al. (2013) listed 15 new potential double eclipsing systems in the SMC. In addition, Čagaš & Pejcha (2012) and Lohr et al. (2013) announced the discovery of new interesting double eclipsing systems. However, a similar study of the period changes for their new discovered systems is still difficult because we lack data spanning a longer time interval.

However, our previous result on the system published in Zasche & Uhlař (2013) seems to be incorrect. The inclination of the eclipsing orbits and their mutual orbit of about 37° is an improbable geometrical configuration and not very stable on longer periods of time. Because of the Kozai cycles (see e.g. Eggleton & Kisseleva-Eggleton 2006), the two orbits should tend to be coplanar or to have a different inclination. And finally, the new minima time observations also significantly deviated from the published predicted fit of V994 Her (for both pairs).

From these reasons we decided to continue monitoring this interesting eclipsing system in the years after our study in 2013 was published. Since then, 36 new observations of minima (of both pairs A and B) were obtained by the authors in seasons 2013, 2014, and 2015 during more than 980 days. The data used for the analysis are presented in Table 1. From studying these new data points, we found that our original hypothesis was incorrect because we had insufficient data. This is well visible in the plot in Fig.1, where we plot $O - C$ diagrams for pairs A and B according to our original solution together with the grey area that represents the time epoch without observations. Unfortunately, this epoch of lacking data is placed exactly in between the two periastron passages in 2006 and 2012. Hence, this paper should not be considered as an erratum, but rather as a reanalysis of the system with new data (new times of minima), as well as a reanalysis of the older radial velocities by Lee et al. (2008). We finally introduce the correct hypothesis of the true system configuration.

3. Updated analysis

The star with the new data was analysed using the same approach as in our previous study. This means that the times of minima were fitted using all 16 parameters for the orbits of the two binaries (the two ephemerides, two apsidal motion hypotheses, and the period changes that are due to the orbit around a common barycenter). The well-known light-travel time effect (LTTE) hypothesis (Irwin, 1959) was used to describe the variation in the $O - C$ diagrams.

In Fig. 2 we plot the updated $O - C$ diagrams for the two A and B pairs, showing their period changes (the y-axis) with respect to time (the x-axis). Our current hypothesis describes the observed data much better (the χ^2 resulted in a value about five times lower). The parameters describing the LTTE fit are given below in Table 2. The orbital period of the two pairs resulted in about half the value presented in our previous study (Zasche & Uhlař, 2013), which is also due to the missing data near the periastron passage in 2009. With the new data the sharp curvatures near the periastron passages are clearly visible and well covered with the observations.

Our hypothesis based on the much shorter orbital period of the two pairs is also able to solve the problem of radial ve-




Fig. 1. Plot of the $O - C$ diagrams of the two pairs A and B. The dots stand for the primary, while the open circles for the secondary minima, the bigger the symbol, the higher the weight. The red dash-dotted lines indicate the LTTE fit, while the black and blue curves represent the final fit (LTTE plus the apsidal motion). This solution is taken from our original study in 2013; the grey area indicates where observations are missing exactly in between the two periastron passages.




Fig. 2. Updated plot of the $O - C$ diagrams of the two pairs A and B (see Fig.1 for description). The missed periastron passage in 2009 is clearly seen in the grey area.

locities published in Lee et al. (2008). The radial velocities they reported are somewhat distorted. Especially those of pair B were unusual because some jumps between the individual datasets near the quadrature are visible. Using our approach, we are also able to compute the radial velocities on the mutual orbit and subtract this contribution from the radial velocities. This is shown in Figure 3, where the calculated radial velocities are shown exactly in the time epochs when the data by Lee et al. (2008) were obtained.

Table 1. New heliocentric minima of V994 Her used for the analysis.

HJD - 2400000	Error	Pair	Type	Filter	Observer/Ref.
56368.61655	0.00029	A	sec	I	RU - this paper
56441.53093	0.00020	A	sec	R	RU - this paper
56463.37125	0.00081	A	prim	C	RU - this paper
56465.45059	0.00259	A	prim	C	RU - this paper
56766.52219	0.00090	A	sec	R	RU - this paper
56767.53147	0.00027	A	prim	R	RU - this paper
56839.44646	0.00247	A	sec	C	RU - this paper
56842.53174	0.00132	A	prim	C	RU - this paper
57119.58372	0.00049	A	prim	C	RU - this paper
57141.49265	0.00127	A	sec	R	RU - this paper
57142.49880	0.00135	A	prim	R	RU - this paper
57189.40527	0.00119	A	sec	R	RU - this paper
57190.41699	0.00066	A	prim	R	RU - this paper
57214.38908	0.00093	A	sec	R	RU - this paper
57215.41792	0.00039	A	prim	R	RU - this paper
57240.41791	0.00029	A	prim	R	RU - this paper
57241.49825	0.00055	A	sec	R	RU - this paper
57287.33273	0.00062	A	sec	C	RU - this paper
57336.25467	0.00038	A	prim	C	RU - this paper
56354.60974	0.00207	B	sec	C	RU - this paper
56450.37231	0.00204	B	prim	C	RU - this paper
56455.43271	0.00040	B	sec	C	RU - this paper
56457.46946	0.00054	B	prim	R	RU - this paper
56736.59690	0.00049	B	sec	R	RU - this paper
56758.50910	0.00259	B	prim	R	RU - this paper
56827.48758	0.00078	B	sec	I	RU - this paper
57091.63291	0.00116	B	sec	R	RU - this paper
57123.50228	0.00057	B	prim	R	RU - this paper
57128.56049	0.00279	B	sec	R	RU - this paper
57133.42997	0.00248	B	prim	R	RU - this paper
57150.47067	0.00049	B	prim	R	RU - this paper
57204.42643	0.00054	B	prim	R	RU - this paper
57209.49597	0.00062	B	sec	I	RU - this paper
57241.34185	0.00229	B	prim	R	RU - this paper
57248.44349	0.00149	B	prim	R	RU - this paper
57295.30457	0.00069	B	prim	R	RU - this paper

The difference between the first and the last data point of about $10 \text{ km} \cdot \text{s}^{-1}$ for pair A and of about $16 \text{ km} \cdot \text{s}^{-1}$ for pair B is significant enough for such a shift. Hence, we applied this correction to the radial velocities by Lee et al. (2008) and reanalysed their data. We found the mass ratios to be only slightly different (the change is below 0.05). We discuss the masses in the next section.

4. Results

Our hypothesis led to several important results, which are completely different from the findings published in the previous analysis. At first, the orbital motion has a period of only about 2.9 years. The other parameters also slightly changed, but the amplitudes and relatively high eccentricity of the mutual orbit remained almost unchanged. However, the apsidal motion of the two pairs resulted in (116 ± 50) years for pair A and (111 ± 40) for pair B. But the most significant change is the resulting geometry of the whole system, which now seems to be close to coplanar (i.e. the orbits of the two eclipsing pairs and their mutual orbit seem to lie very close to one plane, see below). And moreover, if we adopt the Hipparcos (Perryman et al., 1997) value of the parallax as correct ($\pi_{HIP} = (3.90 \pm 0.74) \text{ mas}$), the predicted angular separation of the components is about $\Delta\alpha_{12} = (16.7 \pm 5.6) \text{ mas}$ (instead of the original $\Delta\alpha_{12} = (27.6 \pm 6.8) \text{ mas}$). This value is still rather high and should encourage a prospective interferometric direct detection of the two pairs in the sky. However, the attempt of detection should be made in specific phases on the 2.9 yr orbit to be successful.




Fig. 3. Radial velocities of V994 Her as re-interpreted from the original paper by Lee et al. (2008). *Top:* Radial velocity corrections of the two A and B pairs as these pairs revolve on the orbit around the common barycenter. *Middle:* Radial velocities of pair A (before (plus and cross signs) and after (dot signs) the correction). *Bottom:* Radial velocities of pair B (the same notation).

As a byproduct of our analysis, we also found that the masses of all components as derived by Lee et al. (2008) need to be reconsidered. The masses of the two pairs were calculated using the fact that a so-called mass function of the third body (see e.g. Mayer 1990) can be computed from the orbital parameters, and the inclinations of the two orbits are also close to 90° , that is,

$$\frac{m_A \cdot A_A}{\sin i_A} = \frac{m_B \cdot A_B}{\sin i_B} \implies m_A \cdot A_A \approx m_B \cdot A_B \doteq 0.050,$$


where the masses of pairs are denoted as m_A , and m_B (in solar masses), respectively, and the amplitudes of the period variations in the $O-C$ diagrams for the two pairs are labelled A_A and A_B (in days). The correct mass values are slightly higher for pair A but remained similar for pair B, see Table 3 for a comparison of Lee et al. (2008) and our results. The plot with the resulting masses for the A and B binaries with respect to the inclination is plotted in Fig. 4. The derived mass of pair A is significantly higher than the value published by Lee et al. (2008). Moreover, the plot

Table 2. Updated orbital parameters for V994 Her.

Parameter	Unit	Pair A		Pair B	
		Value	Error	Value	Error
JD_0	HJD	2453855.1339	0.0034	2453857.4361	0.0054
P	Day	2.0832649	0.0000029	1.4200401	0.0000042
e		0.0307	0.0085	0.1253	0.0069
ω	Deg	9.9	6.0	300.7	4.9
$d\omega/dt$	Deg/Cycle	0.0018	0.0012	0.0126	0.0073
p_3	Year	2.91	0.14	2.91	0.14
T_0	HJD	2456045	122	2456045	122
A	Day	0.0091	0.0032	0.0134	0.0037
ω_3	Deg	236.2	26.7	56.2	26.7
e_3		0.758	0.071	0.758	0.071

Table 3. V994 Her: Updated component masses.

	Lee et al. (2008)	This paper
M_{Aa} [M_\odot]	2.83 ± 0.20	3.01 ± 0.06
M_{Ab} [M_\odot]	2.30 ± 0.16	2.58 ± 0.05
M_{Ba} [M_\odot]	1.87 ± 0.12	1.84 ± 0.03
M_{Bb} [M_\odot]	1.86 ± 0.12	1.93 ± 0.04


Fig. 4. Plot of the masses (total mass of the pair A and B) as derived from the analysis of period changes with respect to the inclination of the 2.9 yr orbit. The red solid lines represent the masses reported by Lee et al. (2008), the blue lines stand for the masses from our analysis (with the grey area showing the uncertainties). The black line represents the different masses with respect to the inclination between the orbits.

shows that the inclination of the wide orbit is probably close to 90° , hence the system is close to coplanar. This means that the mutual inclination angle is $i \approx (84 \pm 10)^\circ$ for pair A and $i \approx (85 \pm 12)^\circ$ for pair B.

The question also arises whether it is possible to detect some orbital precession when the two orbits are close to coplanar, and whether we can see some mutual eclipses

of all components in the future. Only a dedicated spectroscopic, photometric, and interferometric monitoring is able to answer this question.

5. Conclusion

The system V994 Her was reanalysed with new photometric observations. Our solution yielded a very different geometrical configuration than originally assumed. In this 2+2 quadruple system with coplanar orbits, the precession of all orbits is only very slow and hard to detect. This system structure is much more probable and seems to be adequately stable for a long period of time. There still remain some open questions, such as the other parameters of the 2.9-year orbit, and the confirmation that the distant component also belongs to the system (which would make this a quintuplet). However, the most promising way to proceed today seems to be an interferometric detection of the two components because they should have a semi-major axis of about 17 mas and a magnitude difference between the components of about 1.4 mag. This is well within the limits for modern stellar interferometers.

The main advantage of the system V994 Her is that it is bright enough for photometric monitoring by only small telescopes (all of our new data were obtained with telescopes with apertures of 20 cm or smaller). It is also quite promising that the periods are relatively short, which means that this is an ideal laboratory of celestial mechanics.

Acknowledgments

We would like to thank David Vokrouhlický for a useful discussion and valuable comments. We also thank the anonymous referee for helpful and critical suggestions. This work was supported by the Czech Science Foundation grants no. P209/10/0715 and GA15-02112S. This research has made use of the Washington Double Star Catalog maintained at the U.S. Naval Observatory, the SIMBAD database, operated at CDS, Strasbourg, France, and of NASA's Astrophysics Data System Bibliographic Services.

References

- Cagaš, P., & Pejcha, O. 2012, A&A, 544, L3
- Duchêne, G., & Kraus, A. 2013, ARA&A, 51, 269
- Eggleton, P. P., & Kiseleva-Eggleton, L. 2006, Ap&SS, 304, 75
- Goodwin, S. P., & Kroupa, P. 2005, A&A, 439, 565
- Goodwin, S. P., Kroupa, P., Goodman, A., & Burkert, A. 2007, Protostars and Planets V, 133
- Heggie, D. C. 1975, MNRAS, 173, 729

- Irwin, J. B. 1959, *AJ*, 64, 149
- Kazarovets, E. V., Samus, N. N., Durlevich, O. V., et al. 1999, *IBVS*, 4659, 1
- Lee, C.-U., Kim, S.-L., Lee, J. W., et al. 2008, *MNRAS*, 389, 1630
- Lohr, M. E., Norton, A. J., Kolb, U. C., et al. 2013, *A&A*, 549, A86
- Mason, B. D., Wycoff, G. L., Hartkopf, W. I., Douglass, G. G., & Worley, C. E. 2001, *AJ*, 122, 3466
- Mayer, P. 1990, *BAICz*, 41, 231
- Orosz, J. A., Welsh, W. F., Carter, J. A., et al. 2012, *Science*, 337, 1511
- Pawlak, M., Graczyk, D., Soszyński, I., et al. 2013, *AcA*, 63, 323
- Perryman, M. A. C., Lindegren, L., Kovalevsky, J., et al. 1997, *A&A*, 323, L49
- Tohline, J. E. 2002, *ARA&A*, 40, 349
- Tokovinin, A. 2008, *MNRAS*, 389, 925
- Zasche, P., & Uhlář, R. 2013, *MNRAS*, 429, 3472